

University of Groningen

Morphological measurements in computed tomography correlate with airflow obstruction in chronic obstructive pulmonary disease

Xie, X.Q.; de Jong, P.A.; Oudkerk, M.; Wang, Y.; ten Hacken, N.H.T.; Miao, J.T.; Zhang, GuiXiang ; de Bock, G.H.; Vliegenthart, R.

Published in:
European Radiology

DOI:
[10.1007/s00330-012-2480-8](https://doi.org/10.1007/s00330-012-2480-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Xie, X. Q., de Jong, P. A., Oudkerk, M., Wang, Y., ten Hacken, N. H. T., Miao, J. T., Zhang, G., de Bock, G. H., & Vliegenthart, R. (2012). Morphological measurements in computed tomography correlate with airflow obstruction in chronic obstructive pulmonary disease: systematic review and meta-analysis. *European Radiology*, 22(10), 2085-2093. <https://doi.org/10.1007/s00330-012-2480-8>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Morphological measurements in computed tomography correlate with airflow obstruction in chronic obstructive pulmonary disease: systematic review and meta-analysis

XueQian Xie · Pim A. de Jong · Matthijs Oudkerk ·
Ying Wang · Nick H. T. ten Hacken · Jingtao Miao ·
GuiXiang Zhang · Geertruida H. de Bock ·
Rozemarijn Vliegenthart

Received: 12 January 2012 / Revised: 20 March 2012 / Accepted: 26 March 2012 / Published online: 15 June 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract

Objectives To determine the correlation between CT measurements of emphysema or peripheral airways and airflow obstruction in chronic obstructive pulmonary disease (COPD).

Methods PubMed, Embase and Web of Knowledge were searched from 1976 to 2011. Two reviewers independently screened 1,763 citations to identify articles that correlated CT measurements to airflow obstruction parameters of the pulmonary function test in COPD patients, rated study quality and extracted information. Three CT measurements were accessed: lung attenuation area percentage < -950 Hounsfield units, mean lung density and airway wall area percentage. Two airflow obstruction parameters were accessed: forced expiratory volume in the first second as percentage

from predicted (FEV_1 %pred) and FEV_1 divided by the forced volume vital capacity.

Results Seventy-nine articles (9,559 participants) were included in the systematic review, demonstrating different methodologies, measurements and CT airflow obstruction correlations. There were 15 high-quality articles (2,095 participants) in the meta-analysis. The absolute pooled correlation coefficients ranged from 0.48 (95 % CI, 0.40 to 0.54) to 0.65 (0.58 to 0.71) for inspiratory CT and 0.64 (0.53 to 0.72) to 0.73 (0.63 to 0.80) for expiratory CT.

Conclusions CT measurements of emphysema or peripheral airways are significantly related to airflow obstruction in COPD patients. CT provides a morphological method to investigate airway obstruction in COPD.

Electronic supplementary material The online version of this article (doi:10.1007/s00330-012-2480-8) contains supplementary material, which is available to authorized users.

X. Xie · M. Oudkerk · R. Vliegenthart (✉)
Center for Medical Imaging–North East Netherlands (CMI-NEN),
Department of Radiology, University of Groningen,
University Medical Center Groningen,
Hanzeplein 1, P.O. Box 30.001, 9700 RB Groningen,
The Netherlands
e-mail: r.vliegenthart@umcg.nl

P. A. de Jong
Department of Radiology, University Medical Center Utrecht,
University of Utrecht,
Heidelberglaan 100, P.O. Box 85.500, 3584 CX Utrecht,
The Netherlands

Y. Wang
Department of Radiology, Tianjin Medical University General
Hospital,
Anshan Road 154,
300052, Tianjin, China

N. H. T. ten Hacken
Department of Pulmonary Diseases, University of Groningen,
University Medical Center Groningen,
Hanzeplein 1, P.O. Box 30.001, 9700 RB Groningen,
The Netherlands

J. Miao · G. Zhang
Department of Radiology, Shanghai Jiao Tong University
Affiliated First People's Hospital,
Haining Road 100,
200080, Shanghai, China

G. H. de Bock
Department of Epidemiology, University of Groningen,
University Medical Center Groningen,
Hanzeplein 1, P.O. Box 30.001, 9700 RB Groningen,
The Netherlands

Key Points

- *Computed tomography is widely performed in patients with chronic obstructive pulmonary disease (COPD)*
- *CT provides quantitative morphological methods to investigate airflow obstruction in COPD*
- *CT measurements correlate significantly with the degree of airflow obstruction in COPD*
- *Expiratory CT measurements correlate more strongly with airflow obstruction than inspiratory CT*
- *Low-dose CT decreases the radiation dose for diagnosis and quantitative emphysema evaluation*

Keywords Review, systematic · Meta-analysis · Tomography, X-ray computed · Pulmonary disease, chronic obstructive · Function test, pulmonary

Abbreviations

%LAA-950	Percentage of lung attenuation area under -950 HU
Ai	Airway lumen area
ATS	American Thoracic Society
CC	Correlation coefficient
CI	Confidence interval
COPD	Chronic obstructive pulmonary disease
CT	Computed tomography
ERS	European Respiratory Society
FEV ₁ %pred	Forced expiratory volume in the first second as percentage from predicted
FEV ₁ /FVC	FEV ₁ divided by the forced volume vital capacity
GOLD	Global initiative for chronic Obstructive Lung Disease
HU	Hounsfield unit
LV	Lung volume
MLD	Mean lung density
Perc15	15 percentile point of lung density
PFT	Pulmonary function test
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-analyses
QUADAS	Quality Assessment of Diagnostic Accuracy Studies
WA%	Wall area percentage
WT	Airway wall thickness

Introduction

Chronic obstructive pulmonary disease (COPD) is characterised by airflow limitation that is not fully reversible [1]. The pathogenesis of airflow limitation in COPD is mainly

related to emphysema and small airway remodelling [2]. Although airflow obstruction parameters in the pulmonary function test (PFT) by spirometry is essential in COPD diagnosis, these parameters fail to quantify the proportionate impact of emphysema and small airways disease individually. Morphological changes can be characterised and quantified by computed tomography (CT), especially by multi-detector CT [3]. For COPD patients, quantitative chest CTs are important for understanding the pathogenesis and the effect of therapeutic interventions [4], and can help to identify those most at risk for acute exacerbations [5]. Since the introduction of the ‘density mask’ in 1988, CT emphysema quantification has had a long history [6–8]. Multi-detector CT can accurately evaluate emphysema [9]. However, quantification of airway remodelling by CT is challenging because of its spatial resolution. Airway wall quantification started over a decade ago, mainly for large airways [3], but investigators have measured the peripheral airways down to 0.5-mm-lumen diameter [10] and 2.8 mm outer diameter [11]. Measurement of narrowing of CT-detectable airways may estimate the degree of small airways disease [10].

The assumption is that emphysema and peripheral airway wall thickness, as detected by CT, are correlated to airflow obstruction in COPD patients. Study results have been variable and sometimes conflicting. However, some individual studies have been small and underpowered [2, 12–15]. Therefore, we conducted a systemic review and meta-analysis to determine the correlation between emphysema or peripheral airway measurements on inspiratory and expiratory CT, and airflow obstruction in COPD.

Materials and methods

This study was conducted according to Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) [16].

Data sources and searches

We searched PubMed, Embase and Web of Knowledge from January 1976 to December 2011, from the start of whole-body CT, using terms related to computed tomography and PFT (i.e. lung function*, respiratory function*, pulmonary function*, etc.) and COPD (i.e. chronic obstructive pulmonary disease*, chronic obstructive lung disease*, etc.) without language restrictions (Electronic supplementary Table 1). Unpublished studies were not included.

Study selection

Four reviewers with at least 6 years' experience in thoracic radiology participated in the study selection. Each study was

evaluated independently by two reviewers out of three, with disagreements resolved by the fourth reviewer. Articles were included in the systematic review if they: (1) analysed the association between CT quantitative emphysema or airway measurements and PFT; (2) investigated human beings; (3) included participants diagnosed with stable adult COPD, according to the Global initiative for chronic Obstructive Lung Disease (GOLD) [1] or the American Thoracic Society (ATS) or the European Respiratory Society (ERS), or clearly defined similar criteria; (4) included participants who had clearly described PFT, according to the guidelines of the ATS, ERS or similar methods. Articles were excluded if they: (1) were reviews, abstracts, case reports or letters; (2) were laboratory or phantom studies; (3) covered participants with confounding disease, such as interstitial lung disease, chronic bronchitis, asthma and α -1 anti-trypsin disease.

Articles were subsequently included in the meta-analysis if they: (1) had no selection bias (e.g. only mild or only severe COPD); (2) had a sample size of ≥ 20 (20 subjects would provide a power of 0.90 when detecting a typical effect correlation coefficient of 0.60); (3) were performed using volumetric multidetector CT; (4) reported correlation coefficients; (5) reported the percentage of lung attenuation area under -950 HU (%LAA-950), mean lung density (MLD) or wall area percentage (WA%) in airways \geq fifth airway generation (sub-sub-segment level) as CT measurements; (6) reported the predicted forced expiratory volume in the first second as percentage (FEV₁ %pred) and FEV₁ divided by the forced volume vital capacity (FEV₁/FVC) as spirometry parameters. Studies were excluded if the CT examination only included selected pulmonary levels or if the slice increment was larger than the slice thickness. In possible duplicate reports, the report with the largest sample size was included.

Data extraction and quality assessment

Two reviewers evaluated independently, with disagreements resolved by a third reviewer. A standardised extraction form was used to collect study characteristics, participant characteristics, methodology and correlation coefficients. The systematic review included ten CT measurements: %LAA-960, %LAA-950, %LAA-910, %LAA-900, MLD, 15 percentile point of lung density (Perc15), lung volume (LV), WA%, airway wall thickness (WT) and airway lumen area (Ai). Three CT measurements, %LAA-950, MLD and WA% were pooled in the meta-analysis. Two PFT parameters for airflow obstruction were collected, including FEV₁ %pred and FEV₁/FVC.

Methodological quality and potential sources of bias of the included meta-analysis articles were assessed with 14 standard items of the Quality Assessment of Diagnostic Accuracy Studies (QUADAS) tool [17]. For each article, a

quality score was accumulated by assigning 1 point to each fulfilled QUADAS item, 0.5 to unclear items and 0 to unmet items. A score of ≥ 11 points was considered high quality and a score < 11 points as low quality. Cohen's k was calculated to indicate inter-observer agreement. Publication bias was evaluated with Begg and Mazumdar rank correlation, Egger's regression test and Rosenthal fail-safe n test.

Data synthesis and analysis

Summary measure was the correlation coefficient (CC). Pooled CCs with 95 % confidence intervals (CIs) were calculated using Hedges-Vervee random effects model and Z-test for normality. Pooled CCs were calculated for the correlations between %LAA-950 and FEV₁ %pred, %LAA-950 and FEV₁/FVC, MLD and FEV₁ %pred, MLD and FEV₁/FVC, and WA% and FEV₁ %pred in inspiratory and expiratory CT. If multi-level bronchi were evaluated, we chose the smallest bronchi. Heterogeneity was tested using the Q statistic and I^2 index. The random effects model was used regardless of the heterogeneity test, although results in the Q statistic were still stated. To investigate the impact of individual variables on the meta-analysis results, subgroup analysis was performed if a subgroup consisted of at least two studies. Subgroups were based on radiation dose (low or normal dose) and breath-hold procedure (inspiratory or expiratory). Meta-regression was performed to investigate the influence of gender, if the male percentage was reported by at least three studies. Statistical analysis was performed using SPSS 18.0 (IBM, NY, USA) and R 2.12.0 (R Foundation, Vienna, Austria). $P < 0.05$ was considered statistically significant.

Results


Study selection

The database searches elicited 1,763 citations (Fig. 1). Seventy-nine articles were included in the systematic review and 15 articles [13, 18–31] in the meta-analysis, including 10 [18–23, 25, 26, 28, 29] from trial cohorts.

Systematic review

The systematic review included 9,559 COPD participants (range of mean age, 48 to 73 years), including 6,101 (63.8 %) men, 2,000 (20.9 %) women and 1,458 (15.3 %) without indicated gender (Electronic supplementary Table 2). A total of 6,935 (72.5 %) were (ex-)smokers and 133 (1.4 %) non-smokers; for 2,491 (26.1 %) no smoking status was reported. Fifty-four (68.4 %) studies were prospective and 23 (29.1 %) retrospective; in 2 (2.5 %) articles the study

Fig. 1 Flowchart of literature review and selection. COPD = Chronic obstruction pulmonary disease; PFT = pulmonary function test; MDCT = multi-detector computed tomography




design was not reported. Forty-one (51.9 %) articles were from Europe, 23 (29.1 %) from Asia and 15 (19.0 %) from North America. Of the articles, 69 (87.3 %) were written in English, 5 (6.3 %) in Italian, 3 (3.8 %) in Chinese, 1 (1.3 %) in French and 1 (1.3 %) in Polish.

The tendency was towards larger sample size and more advanced CT equipment in recent publications (Fig. 2). Before 2007, only articles with sample size <100 were found. After 2007, larger sample size articles were published. Although

single-slice CT (32 articles, 40.5 %) was continuously used from 1993 to 2009, multi-detector CT has been popular in recent years. Since 2002, 4- and 8-slice multi-detector CT (16 articles, 20.2 %) has been used. Since 2005, 16- and 64-slice multi-detector CT (29 articles, 36.7 %) has been used.

Included articles varied in methodology. Volume acquisition was used in 36 articles (45.6 %) and non-volume acquisition in 43 (54.4 %). Selected slices were acquired in 21 studies (26.6 %) and whole lung in 58 (73.4 %). Median slice

Fig. 2 Sample size of the articles included in the systematic review by year of publication and CT generation. MDCT = Multi-detector computed tomography



thickness was 1.0 mm (range, 0.625–10 mm). Median slice increment was 2.0 mm (range, 0.625–20 mm). Low radiation dose was used in 12 articles (15.2 %) and normal dose in 67 (84.7 %). Both inspiratory and expiratory CT findings were evaluated in 24 articles (30.4 %), and only inspiratory in 54 articles (68.3 %) and expiratory in 1 article (1.3 %).

Sixty-four different CT measurements and 27 different PFT parameters were reported (Electronic supplementary Figs. 4 and 5). Common CT measurements were %LAA-950 in 36 (45.6 %) studies, MLD in 22 (27.8 %), WA% in 17 (21.5 %) and visual score in 17 (21.5 %). Common PFT parameters were FEV₁ %pred in 72 articles (91.1 %) and FEV₁/FVC in 64 articles (81.0 %). Common lung parenchyma thresholds defining emphysema ranged from -900 HU to -960 HU, with the most commonly used threshold being -950 HU. In some studies, different correlations to airflow obstruction parameters in PFT were found with these differing thresholds in the same sample [12, 14, 27, 32]. The CC between %LAA-950 and FEV₁ %pred ranged from -0.67 to -0.09 [12, 13], between MLD and FEV₁ %pred from 0.18 to 0.85 [12, 33], between WA% and FEV₁ %pred from -0.713 to -0.044 [22, 34], between %LAA-950 and FEV₁/FVC -0.75 to -0.09 [12, 35], and between MLD and FEV₁/FVC from 0.21 to 0.89 [12, 33]. In four articles, the CC between Perc15 and FEV₁ %pred in inspiration ranged from 0.09 to 0.62, and the CC between Perc15 and FEV₁/FVC in inspiration from 0.12 to 0.62 [12, 13, 36, 37] (Electronic supplementary Table 3).

Risk of bias in the meta-analysis

All articles included in the meta-analysis were high quality (Electronic supplementary Table 4 and Electronic supplementary Fig. 6). The quality score ranged from 12.5 to 13.5. Suboptimal scores were present for three QUADAS items: 7 articles without an interval between CT and PFT (item 4), 9 articles without an indication whether CT quantification was blinded to PFT (item 10) and 13 articles without an indication whether PFT was blinded to CT quantification (item 11). Cohen's κ was 0.925, expressing very good inter-observer agreement. No publication bias was found (Electronic supplementary Table 5). The median of Rosenthal fail-safe n was 122 (range, 84 to 614), indicating a solid empirical result.

Three of the nine meta-analysis calculations showed mild heterogeneity. The I^2 index was >50 % for the correlation between MLD and FEV₁ %pred in inspiration ($P=0.11$, I^2 index=50.6 %), between MLD and FEV₁/FVC in inspiration ($P=0.02$, I^2 index=75.9 %), and between WA% and FEV₁ %pred ($P=0.04$, I^2 index=64.7 %).

Synthesis of results in the meta-analysis

The meta-analysis included 2,095 participants out of 9,559 in the systematic review (Fig.3).

Nine articles [13, 18, 19, 22–24, 26–28, 31] reported CC between %LAA-950 and FEV₁ %pred in inspiration. Two [19, 25] were from the National Lung Screening Trial (NLST) cohort and two [22, 23] from the Korean Obstructive Lung Disease (KOLD) cohort. Because of possible duplicate reporting, articles [22, 25] with smaller sample sizes were excluded from each cohort. The pooled CC between %LAA-950 and FEV₁ %pred was -0.48 (95% CI: -0.54, -0.40) in inspiration and -0.64 (-0.72, -0.53) in expiration. The pooled CC between %LAA-950 and FEV₁/FVC was -0.65 (-0.71, -0.58) in inspiration and -0.69 (-0.78, -0.59) in expiration.

No potential duplicate report was found for MLD. The pooled CC between MLD and FEV₁ %pred was 0.51 (95% CI: 0.38, 0.63) in inspiration and 0.71 (0.58, 0.81) in expiration. The pooled CC between MLD and FEV₁/FVC was 0.64 (0.52, 0.74) in inspiration and 0.73 (0.63, 0.80) in expiration.


Seven articles [18, 20–22, 25, 28, 30] reported CC between WA% and FEV₁ %pred in inspiration. Two articles [18, 22] were excluded because airway measurements concerned only airways above the fifth airway generation. Another article [25] was excluded because it did not report which airways were measured. Four articles [20, 21, 28, 30] were finally included. The lumen diameter of peripheral airways was about 2–3 mm in the included articles. The pooled CC between WA% and FEV₁ %pred was -0.59 (95% CI: -0.68, -0.48) in inspiration. Expiratory CT was not used for airway measurements.

Subgroup analysis and meta-regression

Subgroup analysis for radiation dose was performed for the association between %LAA-950 and FEV₁ %pred in inspiration, indicating no significant difference ($P>0.05$). In low dose [18, 19], the pooled CC was -0.44 (95% CI: -0.50, -0.37). In normal dose [13, 23, 24, 26, 27, 31], the pooled CC was -0.50 (-0.57, -0.42). Subgroup analysis was also performed for inspiratory and expiratory CT. A significantly stronger negative correlation was found between %LAA-950 and FEV₁ %pred in expiratory CT ($P<0.05$), and a stronger positive correlation between MLD and FEV₁ %pred ($P<0.001$), but no difference was found in the association between %LAA-950 and FEV₁/FVC ($P>0.05$), or MLD and FEV₁/FVC ($P>0.05$). In meta-regression for gender contribution, no statistically significant effect modification was found for male percentage ($P>0.05$) (Electronic supplementary Table 5).

Discussion

In this meta-analysis, significant correlations were found between CT measurements of emphysema or peripheral airway and airflow obstruction parameters in PFT in COPD



*: Subgroups analysis was performed between inspiratory and expiratory CT

Fig. 3 Forest plots for correlations between CT measurements and airflow obstruction. CI = Confidence interval; $P(Z) = P$ value of Z test; FEV₁ %pred = percentage of the predicted forced expiratory volume in the first second; FEV₁/FVC = FEV₁ divided by forced vital capacity; %LAA-950 = percentage lower attenuation area than -950 HU; MLD = mean lung density; Perc15 = 15 percentile point of lung density; WA% = wall area percentage

patients, both in inspiratory and expiratory CT. The range of absolute correlation coefficients between included CT measurements and airflow obstruction was 0.48 to 0.65 for inspiratory CT and 0.64 to 0.73 for expiratory CT. These results confirm correlations between morphology and function in COPD patients. The confidence in these findings is strong, as results were based on high methodological quality studies without publication bias. Thus, CT provides a quantitative morphological method to investigate the principle components of airway obstruction in COPD, with similar strength of associations with airflow obstruction for CT measurements of emphysema and peripheral airways. The strongest association was found between CT emphysema measurements and FEV₁/FVC, especially in expiratory CT. Our systematic review demonstrated differing methodologies for CT quantification and contrasting correlations with airflow obstruction in COPD patients.

CT quantification reflects pathophysiological changes in COPD to some degree. The pathological findings of airway limitation are in airways <2 mm in internal diameter [38]. Such small airways can hardly be measured directly by CT because of the spatial resolution limit. However, peripheral airway ($\geq 5^{\text{th}}$ generation) wall thickness can be measured as WA%. Destruction of the lung parenchyma (emphysema) can be measured as %LAA-950 or MLD. The morphological contribution from these two pathological processes is difficult to distinguish by spirometry, but is important for COPD research. The morphological information in CT quantification of the relative predominance of peripheral airway wall disease or emphysema may in the future allow more focused treatment of the predominating COPD phenotype.

This systematic review incorporated ten different CT measurements. Although a visual score was common in earlier publications, we did not discuss it, because of its subjective nature. Perc15 seems an effective measurement for emphysema, but Perc15 results could not be pooled because of an insufficient study number. Only three (%LAA-950, MLD and WA%) of the ten measurements were eventually pooled because the study number was sufficient to perform a meta-analysis. We investigated two PFT parameters (FEV₁ %pred and FEV₁/FVC) as they are the two most commonly used functional parameters regarding airway limitation.

In subgroup analysis, associations for inspiratory and expiratory CT findings were compared. Some authors have reported

that CT measurements in expiration are more closely correlated with airflow obstruction than in inspiration [14, 39–41]. Our results indicate that CT measurements in expiration rather than in inspiration were more correlated with FEV₁ %pred, not with FEV₁/FVC. Whether an additional expiratory CT data acquisition should be performed for COPD evaluation is debatable. Expiratory CT exposes patients to additional radiation; however, with developments in CT technique, the additional dose will likely decrease. Also, we found low radiation doses did not change correlations between CT emphysema quantification and airflow obstruction compared to normal doses. Low-dose CT can decrease the overall radiation dose for CT quantitative emphysema evaluation without loss in diagnostic value.

Multiple airway generations were included in the systematic review, but only peripheral airways ($\geq 5^{\text{th}}$ generation) in the meta-analysis. Some authors investigated airways from the third to fifth or sixth generation, and found that the association between airway wall measurements and PFT was stronger for higher generations than lower generations [20, 28]. Therefore, we only pooled results for airways \geq fifth generation. Some authors reported moderate associations with larger airways, ranging from -0.39 to -0.54 [18, 42]. In our meta-analysis, the association between wall area percentage of peripheral airways and FEV₁ %pred was -0.59. One factor to keep in mind is the overestimation of airway wall thickness, showing a relative increase with each airway generation [3]. Despite this factor, based on our pooled results, the association between disease of the more peripheral airways ($\geq 5^{\text{th}}$ generation) and lung function appears stronger than for lower generation airways ($< 5^{\text{th}}$ generation), suggesting that airway wall thickness measurements on CT should be performed on the smallest airways visible.

This study confirms significant correlations between CT measurements and airflow obstruction in COPD. The correlations were in agreement with some expert narrative reviews [9, 10] and individual studies [14, 15, 39, 42, 43]. Nevertheless, other studies reported weaker associations, e.g. the National Emphysema Treatment Trial (NETT) study [12] and the International COPD Genetics Network (ICGN) study [2]. In NETT and ICGN, predominantly single-slice CT was used. Since single-slice CT decreases reproducibility and accuracy [44], this has likely caused the reduced strength of the correlations.

This study has some limitations. First, no prospective large cohort with up-to-date CT technology was found as primary study. The largest study [45] included over 2,000 participants, but most of the participants had normal lung function. In inspiratory CT, the number of included articles was relatively small, but sufficient to perform a reliable meta-analysis. However, in expiratory CT, some indicators of bias could not be determined because of the limited number of studies. Second, we found 64 different quantitative CT measurements in the

literature. We chose ten for systematic review and three for meta-analysis. Although the included parameters are representative, the other 54 measurements could be valuable to evaluate COPD. In another way, FEV₁ %pred and FEV₁/FVC were selected as airflow obstruction parameters in PFT, but other parameters in PFT were valuable for evaluating COPD. Third, mild heterogeneity was found in three correlations in the meta-analysis. A random effects model was used to compensate for the heterogeneity.

In conclusion, measurements of emphysema and the peripheral airways on inspiratory and expiratory CT have significant correlations with airflow obstruction as assessed by FEV₁ %pred and FEV₁/FVC in COPD patients. Thus, CT provides a quantitative morphological method to investigate airflow obstruction by emphysema and peripheral airway disease in COPD.

Acknowledgments We thank Dr. Estelle J.K. Noach, Dr. Peter M.A. van Ooijen and Mrs. Y. Zhao, all from the Center for Medical Imaging–North East Netherlands (CMI-NEN), Department of Radiology of the University of Groningen, University Medical Center Groningen, for revision of the manuscript, methodological guidance and study selection, respectively.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Rabe KF, Hurd S, Anzueto A et al (2007) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease—GOLD executive summary. *Am J Respir Crit Care Med* 176:532–555
2. Patel BD, Coxson HO, Pillai SG et al (2008) Airway wall thickening and emphysema show independent familial aggregation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 178:500–505
3. Nakano Y, Muro S, Sakai H et al (2000) Computed tomographic measurements of airway dimensions and emphysema in smokers correlation with lung function. *Am J Respir Crit Care Med* 162:1102–1108
4. Coxson HO (2008) Quantitative chest tomography in COPD research: chairman's summary. *Proc Am Thorac Soc* 5:874–877
5. Han MK, Kazerooni EA, Lynch DA et al (2011) Chronic obstructive pulmonary disease exacerbations in the COPDGene study: associated radiologic phenotypes. *Radiology* 261:274–282
6. Muller NL, Staples CA, Miller RR, Abboud RT (1988) "Density mask". An objective method to quantitate emphysema using computed tomography. *Chest* 94:782–787
7. Guenard H, Diallo MHH, Laurent F, Vergeret J (1992) Lung density and lung mass in emphysema. *Chest* 102:198–203
8. Lamers RJ, Thelissen GR, Kessels AG, Wouters EF, van Engelshoven JM (1994) Chronic obstructive pulmonary disease: evaluation with spirometrically controlled CT lung densitometry. *Radiology* 193:109–113
9. Matsuoka S, Yamashiro T, Washko GR, Kurihara Y, Nakajima Y, Hatabu H (2010) Quantitative CT assessment of chronic obstructive pulmonary disease. *Radiographics* 30:55–66
10. de Jong PA, Muller NL, Pare PD, Coxson HO (2005) Computed tomographic imaging of the airways: relationship to structure and function. *Eur Respir J* 26:140–152
11. Kosciuch J, Krenke R, Gorska K, Zukowska M, Maskey-Warzechowska M, Chazan R (2009) Relationship between airway wall thickness assessed by high-resolution computed tomography and lung function in patients with asthma and chronic obstructive pulmonary disease. *J Physiol Pharmacol* 60:71–76
12. Washko GR, Criner GJ, Mohsenifar Z et al (2008) Computed tomographic-based quantification of emphysema and correlation to pulmonary function and mechanics. *COPD* 5:177–186
13. Akira M, Toyokawa K, Inoue Y, Arai T (2009) Quantitative CT in chronic obstructive pulmonary disease: inspiratory and expiratory assessment. *Am J Roentgenol* 192:267–272
14. Matsuoka S, Kurihara Y, Yagihashi K, Nakajima Y (2007) Quantitative assessment of peripheral airway obstruction on paired expiratory/inspiratory thin-section computed tomography in chronic obstructive pulmonary disease with emphysema. *J Comput Assist Tomogr* 31:384–389
15. Orlandi I, Moroni C, Camiciottoli G et al (2004) Spirometric-gated computed tomography quantitative evaluation of lung emphysema in chronic obstructive pulmonary disease—a comparison of 3 techniques. *J Comput Assist Tomogr* 28:437–442
16. Moher D, Liberati A, Tetzlaff J, Altman DG (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Br Med J* 339:332–336
17. Whiting PF, Weswood ME, Rutjes AW, Reitsma JB, Bossuyt PN, Kleijnen J (2006) Evaluation of QUADAS, a tool for the quality assessment of diagnostic accuracy studies. *BMC Med Res Methodol* 6:9
18. Bon JM, Leader JK, Weissfeld JL et al (2009) The influence of radiographic phenotype and smoking status on peripheral blood biomarker patterns in chronic obstructive pulmonary disease. *PLoS One* 4:e6865
19. Dransfield MT, Washko GR, Foreman MG, Estepar RSJ, Reilly J, Bailey WC (2007) Gender differences in the severity of CT emphysema in COPD. *Chest* 132:464–470
20. Hasegawa M, Nasuhara Y, Onodera Y et al (2006) Airflow limitation and airway dimensions in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 173:1309–1315
21. Leader JK, Zheng B, Sciruba FC et al (2008) The relation of airway size to lung function. *Proc SPIE Med Imaging* 6916:691623
22. Lee YK, Oh YM, Lee JH et al (2008) Quantitative assessment of emphysema, air trapping, and airway thickening on computed tomography. *Lung* 186:157–165
23. Park YS, Seo JB, Kim N et al (2008) Texture-based quantification of pulmonary emphysema on high-resolution computed tomography: Comparison with density-based quantification and correlation with pulmonary function test. *Investig Radiol* 43:395–402
24. Pauls S, Gulkin D, Feuerlein S et al (2010) Assessment of COPD severity by computed tomography: correlation with lung functional testing. *Clin Imaging* 34:172–178
25. Washko GR, Dransfield MT, Estepar RS et al (2009) Airway wall attenuation: a biomarker of airway disease in subjects with COPD. *J Appl Physiol* 107:185–191
26. Yamashiro T, Matsuoka S, Bartholmai BJ et al (2010) Collapsibility of lung volume by paired inspiratory and expiratory CT scans. Correlations with lung function and mean lung density. *Acad Radiol* 17:489–495
27. Zhang J, Yu TL, Wu Q, Ye N, Cao Y, Zhou N (2008) Quantitative MSCT study of lung attenuation in patients with chronic obstructive pulmonary disease. *Chin J Med Imaging Technol* 24:1611–1614

28. Yamashiro T, Matsuoka S, San Jose Estepar R et al (2010) Quantitative assessment of bronchial wall attenuation with thin-section CT: an indicator of airflow limitation in chronic obstructive pulmonary disease. *Am J Roentgenol* 195:363–369
29. Hesselbacher SE, Ross R, Schabath MB et al (2011) Cross-sectional analysis of the utility of pulmonary function tests in predicting emphysema in ever-smokers. *Int J Environ Res Public Health* 8:1324–1340
30. Ohno Y, Koyama H, Yoshikawa T et al (2011) Comparison of capability of dynamic O2-enhanced MRI and quantitative thin-section MDCT to assess COPD in smokers. *Eur J Radiol*. doi:10.1016/j.ejrad.2011.1002.1004
31. Iwasawa T, Takahashi H, Ogura T et al (2011) Influence of the distribution of emphysema on diaphragmatic motion in patients with chronic obstructive pulmonary disease. *Jpn J Radiol* 29:256–264
32. Jin LF, Miao JT, Li ZY, Xie XQ, Dai XM (2007) Study of voxel index from multislice CT in evaluating chronic obstructive pulmonary disease air-trapping. *Chin J Med Imaging Technol* 23:1033–1036
33. Falaschi F, Miniati M, Battolla L et al (1995) Quantification of pulmonary emphysema with computerized tomography. Comparison with various methods. *Radiol Med* 90:16–23
34. Deveci F, Murat A, Turgut T, Altuntas E, Muz MH (2004) Airway wall thickness in patients with COPD and healthy current smokers and healthy non-smokers: assessment with high resolution computed tomographic scanning. *Respiration* 71:602–610
35. Iwasawa T, Takahashi H, Ogura T et al (2007) Correlation of lung parenchymal MR signal intensity with pulmonary function tests and quantitative computed tomography (CT) evaluation: a pilot study. *J Magn Reson Imaging* 26:1530–1536
36. Heussel CP, Herth FJ, Kappes J et al (2009) Fully automatic quantitative assessment of emphysema in computed tomography: comparison with pulmonary function testing and normal values. *Eur Radiol* 19:2391–2402
37. Shaker SB, Maltbaek N, Brand P, Haeussermann S, Dirksen A (2005) Quantitative computed tomography and aerosol morphometry in COPD and alpha-antitrypsin deficiency. *Eur Respir J* 25:23–30
38. Hogg JC, Chu F, Utokaparch S et al (2004) The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 350:2645–2653
39. Camiciottoli G, Bartolucci M, Maluccio NM et al (2006) Spirometrically gated high-resolution CT findings in COPD—lung attenuation vs lung function and dyspnea severity. *Chest* 129:558–564
40. O'Donnell RA, Peebles C, Ward JA et al (2004) Relationship between peripheral airway dysfunction, airway obstruction, and neutrophilic inflammation in COPD. *Thorax* 59:837–842
41. Zaporozhan J, Ley S, Eberhardt R et al (2005) Paired inspiratory/expiratory volumetric thin-slice CT scan for emphysema analysis: comparison of different quantitative evaluations and pulmonary function test. *Chest* 128:3212–3220
42. Achenbach T, Weinheimer O, Biedermann A et al (2008) MDCT assessment of airway wall thickness in COPD patients using a new method: correlations with pulmonary function tests. *Eur Radiol* 18:2731–2738
43. Boschetto P, Quintavalle S, Zeni E et al (2006) Association between markers of emphysema and more severe chronic obstructive pulmonary disease. *Thorax* 61:1037–1042
44. Newell JD Jr (2008) Quantitative computed tomography of lung parenchyma in chronic obstructive pulmonary disease: an overview. *Proc Am Thorac Soc* 5:915–918
45. Mohamed Hoesein FAA, De Hoop B, Zanen P et al (2011) CT-quantified emphysema in male heavy smokers: association with lung function decline. *Thorax* 66:782–787